

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.079$
 $S = 1.121$
 3077 reflections
 239 parameters
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0452P)^2 + 0.3435P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.038$
 $\Delta\rho_{\max} = 0.439 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.518 \text{ e } \text{Å}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick, 1997a)
 Extinction coefficient: 0.0948 (9)
 Scattering factors from *International Tables for Crystallography* (Vol. C)
 Absolute structure: Flack (1983)
 Flack parameter = -0.03 (3)

Table 1. Selected geometric parameters (Å , $^\circ$)

Sb1—Cl1	2.738 (1)	Sb1—Cl4	2.768 (1)
Sb1—Cl2	2.504 (1)	Sb1—Cl5	2.371 (1)
Sb1—Cl3	2.499 (1)	Sb1...Cl5 ⁱ	3.474 (1)
Cl1—Sb1—Cl2	174.90 (4)	Cl2—Sb1—Cl4	90.19 (4)
Cl1—Sb1—Cl3	89.60 (4)	Cl2—Sb1—Cl5	89.12 (4)
Cl1—Sb1—Cl4	89.67 (4)	Cl3—Sb1—Cl4	178.60 (4)
Cl1—Sb1—Cl5	85.78 (4)	Cl3—Sb1—Cl5	90.41 (4)
Cl2—Sb1—Cl3	90.44 (4)	Cl4—Sb1—Cl5	88.35 (4)

Symmetry code: (i) $1 + x, y, z$.Table 2. Hydrogen-bonding geometry (Å , $^\circ$)

D—H...A	D—H	H...A	D...A	D—H...A
N11—H11D...Cl1	0.90	2.35	3.238 (9)	170
N12—H12D...Cl1	0.90	2.17	3.054 (8)	169
N21—H21D...Cl4	0.90	2.22	3.114 (10)	176
N22—H22D...Cl4	0.90	2.23	3.132 (11)	177

Data collection: *Kuma Diffraction Software* (Kuma, 1996).
 Cell refinement: *Kuma Diffraction Software*. Data reduction:
Kuma Diffraction Software. Program(s) used to solve structure:
SHELXS97 (Sheldrick, 1997b). Program(s) used to refine
 structure: *SHELXL97* (Sheldrick, 1997a). Molecular graphics:
SHELXTL (Sheldrick, 1990). Software used to prepare material
 for publication: *SHELXL97*.

This work was financially supported by KBN under project No. 3TO9A 104 17.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: KA1330). Services for accessing these data are described at the back of the journal.

References

- Bujak, M. & Zaleski, J. (1998). *Acta Cryst.* **C54**, 1773–1777.
 Ciapała, P., Zaleski, J., Bator, G., Jakubas, R. & Pietraszko, A. (1996). *J. Phys. Condens. Matter*, **8**, 1957–1970.
 Ensinger, U., Schwarz, W. & Schmidt, A. (1983). *Z. Naturforsch. Teil B*, **38**, 149–154.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Jakubas, R., Czapała, Z., Galewski, Z., Sobczyk, L., Żogała, O. J. & Lis, T. (1986). *Phys. Status Solidi A*, **93**, 449–455.
 Knödler, F., Ensinger, U., Schwarz, W. & Schmidt, A. (1988). *Z. Anorg. Allg. Chem.* **557**, 208–218.
 Kuma (1996). *Kuma Diffraction Software*. Versions 8.1.0 and 8.1.1. Kuma Diffraction, Wrocław, Poland.
 Lipka, A. (1980). *Z. Anorg. Allg. Chem.* **469**, 229–233.
 Sheldrick, G. M. (1990). *SHELXTL*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

- Sheldrick, G. M. (1997a). *SHELXL97. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
 Sheldrick, G. M. (1997b). *SHELXS97. Program for the Solution of Crystal Structures*. University of Göttingen, Germany.
 Sobczyk, L., Jakubas, R. & Zaleski, J. (1997). *Pol. J. Chem.* **71**, 265–300.
 Zaleski, J. & Pietraszko, A. (1995). *J. Phys. Chem. Solids*, **56**, 883–890.
 Zaleski, J. & Pietraszko, A. (1996). *Acta Cryst.* **B52**, 287–295.

Acta Cryst. (1999). **C55**, 1778–1780

[*N*-Acetyl-*N'*-methylureato(2-)-*N,N'*](η^5 -pentamethylcyclopentadienyl)(triphenylphosphine-*P*)iridium(III) chloroform solvate

MAARTEN B. DINGER,^a WILLIAM HENDERSON,^a ALLEN G. OLIVER^b AND CLIFTON E. F. RICKARD^b

^aDepartment of Chemistry, University of Waikato, Private Bag 3105, Hamilton, New Zealand, and ^bDepartment of Chemistry, University of Auckland, Private Bag 92019, Auckland, New Zealand. E-mail: w.henderson@waikato.ac.nz

(Received 11 February 1999; accepted 3 June 1999)


Abstract

The title complex, $[\text{Ir}(\text{C}_{10}\text{H}_{15})(\text{C}_4\text{H}_6\text{N}_2\text{O}_2)(\text{C}_{18}\text{H}_{15}\text{P})]\text{-CHCl}_3$, displays a 'piano-stool'-type configuration of urea ligands, with the Ir atom in an Ir—NR—C(O)—NR' ring system. This is the first structure determination of a coordinatively saturated iridaureylene complex.

Comment

Ureylene complexes containing the $M\text{-NR-C(O)-NR}'$ four-membered ring system with a coordinated urea dianion are known for the majority of transition metals (Dinger & Henderson, 1998a; Braunstein & Nobel, 1989; Cenini & La Monica, 1976). A variety of synthetic routes have been described for the preparation of such complexes (Dinger & Henderson, 1998a) and we have recently been investigating the use of silver(I) oxide as a reagent for the synthesis of transition metal ureylene complexes (Dinger & Henderson, 1998a,b; Dinger *et al.*, 1996, 1998). As part of these studies, the iridium(III) ureylene complex $[\text{Cp}^*\text{Ir}\{\text{N}(\text{Ac})\text{C}(\text{O})\text{N}(\text{Ac})\}(\text{PPh}_3)]$ (Cp^* is pentamethylcyclopentadienyl), derived from *N,N'*-diacetylurea, was reported (Dinger *et al.*, 1998). We now report the crystal structure of the analogous iridium(III) complex derived from *N*-acetyl-*N'*-methylurea, $[\text{Cp}^*\text{Ir}\{\text{N}(\text{Ac})\text{C}(\text{O})\text{NMe}\}(\text{PPh}_3)]\text{-CHCl}_3$, (I); the platinum com-

plex derived from this urea has been reported previously (Dinger *et al.*, 1996).



Only one crystal structure of a rhodium or iridium ureylene complex has been described to date, that of the coordinatively unsaturated $[\text{Cp}^*\text{Ir}\{\text{N}(2,6\text{-}^i\text{Pr}_2\text{C}_6\text{H}_3)\text{-C}(\text{O})\text{N}(2,4,6\text{-Me}_3\text{C}_6\text{H}_2)\}]$ (Danopoulos *et al.*, 1996). The accuracy of this structure was rather low.

The title complex (Fig. 1) shows the expected pseudo-six-coordinate structure, with the Cp^* ligand occupying three coordination sites on the iridium centre, which forms part of a four-membered ureylene ring. The Ir—N bond lengths [2.079 (3) and 2.092 (3) Å] are the same within experimental error, though the Ir—N2 bond is tentatively shorter, as might be expected due to strengthening of the Ir—N bond by the electronegative acetyl substituent. Both are rather longer than the Ir—N bond lengths in the complex $[\text{Cp}^*\text{Ir}\{\text{N}(2,6\text{-}^i\text{Pr}_2\text{C}_6\text{H}_3)\text{C}(\text{O})\text{N}(2,4,6\text{-Me}_3\text{C}_6\text{H}_2)\}]$ [1.92 (2) and 2.03 (2) Å; Danopoulos *et al.*, 1996], presumably due to the coordinatively unsaturated nature of this complex. The ureylene C1—O1 bond of the title complex [1.250 (5) Å] is considerably longer than in $[\text{Cp}^*\text{Ir}\{\text{N}(2,6\text{-}^i\text{Pr}_2\text{C}_6\text{H}_3)\text{C}(\text{O})\text{N}(2,4,6\text{-Me}_3\text{C}_6\text{H}_2)\}]$ [1.18 (2) Å], but is comparable with the acetyl C3—O2 bond length [1.236 (5) Å]. The two N—C bond lengths in the iridacycle are unequal in length [N1—C1 1.348 (5) and N2—C1 1.413 (5) Å], the dif-

ference presumably arising from the differing electronegativities of the two nitrogen substituents (acetyl versus methyl) on the ring system. The acetyl substituent lies approximately in the plane of the metallacycle, with the C1—N2—C3—C4 torsion angle being 8.9 (7)°. The Cp^* ligand appears to be reasonably symmetrically bonded, with Ir—C bond lengths in the range 2.210 (3)—2.223 (4) Å. The Ir—P distance of 2.3001 (9) Å is normal for an iridium to phosphine bond.

Experimental

The title complex was prepared by the reaction of $[\text{Cp}^*\text{IrCl}_2(\text{PPh}_3)]$ (Booth *et al.*, 1969) with *N*-acetyl-*N'*-methylurea and excess silver(I) oxide in refluxing dichloromethane, under a nitrogen atmosphere, following a method described previously for this type of complex (Dinger *et al.*, 1996, 1998). The silver salts were removed by filtration and the filtrate evaporated to dryness to give the crude product. Yellow crystals of the title complex as a CHCl_3 solvate were grown by slow evaporation from a chloroform-diethyl ether solution. Electrospray MS (positive-ion mode, 1:1 $\text{MeCN}/\text{H}_2\text{O}$, cone voltage 20 V): m/z 705 $[\text{M} + \text{H}]^+$ (100%).

Crystal data

$[\text{Ir}(\text{C}_{10}\text{H}_{15})(\text{C}_4\text{H}_6\text{N}_2\text{O}_2)\text{-}(\text{C}_{18}\text{H}_{15}\text{P})]\cdot\text{CHCl}_3$
 $M_r = 823.17$
 Monoclinic
 $P2_1/n$
 $a = 12.5471$ (2) Å
 $b = 17.0445$ (1) Å
 $c = 16.7432$ (3) Å
 $\beta = 111.060$ (1)°
 $V = 3341.51$ (8) Å³
 $Z = 4$
 $D_x = 1.636$ Mg m⁻³
 D_m not measured

Mo $K\alpha$ radiation
 $\lambda = 0.71073$ Å
 Cell parameters from 6915 reflections
 $\theta = 1.76\text{--}27.43^\circ$
 $\mu = 4.316$ mm⁻¹
 $T = 203$ (2) K
 Tablet
 $0.39 \times 0.36 \times 0.12$ mm
 Yellow

Data collection

Siemens SMART CCD diffractometer
 Multi-scan
 Absorption correction: empirical (Blessing, 1995)
 $T_{\min} = 0.284$, $T_{\max} = 0.626$
 19 310 measured reflections
 7308 independent reflections

6154 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$
 $\theta_{\max} = 27.43^\circ$
 $h = -15 \rightarrow 14$
 $k = 0 \rightarrow 22$
 $l = 0 \rightarrow 21$
 Intensity decay: none

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.066$
 $S = 1.067$
 7308 reflections
 386 parameters
 H atoms constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0215P)^2 + 5.8390P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.033$
 $\Delta\rho_{\max} = 1.548$ e Å⁻³
 $\Delta\rho_{\min} = -1.039$ e Å⁻³
 Extinction correction: none
 Scattering factors from *International Tables for Crystallography* (Vol. C)




Fig. 1. The molecular structure of the title compound showing 50% probability displacement ellipsoids. The chloroform of crystallization and the H atoms have been omitted for clarity.

Table 1. Selected geometric parameters and hydrogen-bonding geometry (Å, °)

Ir—N1	2.092 (3)	P—C31	1.832 (4)	
Ir—N2	2.079 (3)	P—C41	1.825 (4)	
Ir—C5	2.223 (4)	N1—C1	1.348 (5)	
Ir—C6	2.211 (4)	N1—C2	1.429 (5)	
Ir—C7	2.222 (3)	N2—C1	1.413 (5)	
Ir—C8	2.210 (3)	N2—C3	1.349 (5)	
Ir—C9	2.216 (4)	O1—C1	1.250 (5)	
Ir—P	2.3001 (9)	O2—C3	1.236 (5)	
P—C21	1.833 (4)	C3—C4	1.512 (6)	
N1—Ir—P	87.84 (9)	C3—N2—C1	129.9 (3)	
N2—Ir—P	88.79 (9)	O1—C1—N1	128.2 (4)	
N2—Ir—N1	62.08 (12)	O1—C1—N2	129.5 (4)	
C1—N1—Ir	98.5 (2)	O2—C3—N2	121.3 (4)	
C2—N1—Ir	137.7 (3)	O2—C3—C4	120.6 (4)	
C1—N2—Ir	97.0 (2)	N1—C1—N2	102.3 (3)	
C3—N2—Ir	132.6 (3)	N2—C3—C4	118.1 (4)	
C1—N1—C2	123.8 (3)			
D—H...A	D—H	H...A	D...A	D—H...A
C50—H50A...O1	0.99	2.01	2.958 (6)	159

A single molecule of chloroform of crystallization was located from residual electron-density maps and was refined anisotropically. A reasonable hydrogen bond exists between the chloroform carbon (C50) and the O1 atom [2.958 (6) Å]. The maximum residual peak lies 1.37 Å from C45 and the minimum peak lies 0.89 Å from the Ir atom.

Data collection: SMART (Siemens, 1994a). Cell refinement: SAINT (Siemens, 1994a). Data reduction: SAINT. Program(s) used to solve structure: SHELXS97 (Sheldrick, 1997a). Program(s) used to refine structure: SHELXL97 (Sheldrick, 1997b). Molecular graphics: SHELXTL (Siemens, 1994b). Software used to prepare material for publication: SHELXL97.

We thank the Universities of Waikato and Auckland, together with the New Zealand Lottery Grants Board, for financial support of this work.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: TA1247). Services for accessing these data are described at the back of the journal.

References

- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
 Booth, B. L., Haszeldine, R. N. & Hill, M. (1969). *J. Organomet. Chem.* **16**, 491–496.
 Braunstein, P. & Nobel, D. (1989). *Chem. Rev.* **89**, 1927–1945.
 Cenini, S. & La Monica, G. (1976). *Inorg. Chim. Acta*, **18**, 279–293.
 Danopoulos, A. A., Wilkinson, G., Sweet, T. K. N. & Hursthouse, M. B. (1996). *J. Chem. Soc. Dalton Trans.* pp. 3771–3778.
 Dinger, M. B. & Henderson, W. (1998a). *J. Organomet. Chem.* **557**, 231–241.
 Dinger, M. B. & Henderson, W. (1998b). *J. Chem. Soc. Dalton Trans.* pp. 1763–1773.
 Dinger, M. B., Henderson, W. & Nicholson, B. K. (1998). *J. Organomet. Chem.* **556**, 75–88.
 Dinger, M. B., Henderson, W., Nicholson, B. K. & Wilkins, A. L. (1996). *J. Organomet. Chem.* **526**, 303–312.
 Sheldrick, G. M. (1997a). SHELXS97. Program for the Solution of Crystal Structures. University of Göttingen, Germany.
 Sheldrick, G. M. (1997b). SHELXL97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany.

Siemens (1994a). SMART and SAINT. Area Detector Control and Integration Software. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Siemens (1994b). SHELXTL. Release 5.03. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Acta Cryst. (1999). **C55**, 1780–1784

Triaqua(2,2'-bipyridyl-*N,N'*)(thiosulfato-*S*)-nickel(II) dihydrate and triaqua(1,10-phenanthroline-*N,N'*)(thiosulfato-*S*)nickel(II) monohydrate

ELEONORA FREIRE,^a SERGIO BAGGIO,^a RICARDO BAGGIO^b AND LEOPOLDO SUESCUN^c

^aUniversidad Nacional de la Patagonia, Sede Puerto Madryn, and CenPat, CONICET, 9120 Puerto Madryn, Chubut, Argentina, ^bDepartamento de Física, Comisión Nacional de Energía Atómica, Buenos Aires, Argentina, and ^cCSSC Lab., Facultad de Química, Universidad de la República, Montevideo, Uruguay. E-mail: baggio@cnea.gov.ar

(Received 24 February 1999; accepted 21 June 1999)

Abstract

In the title compounds, [Ni(S₂O₃)(C₁₀H₈N₂)(H₂O)₃]-2H₂O and [Ni(S₂O₃)(C₁₂H₈N₂)(H₂O)₃]-H₂O, the Ni^{II} ion displays a distorted octahedral coordination provided by the two N atoms of the organic ligand, one S atom from the thiosulfate anion and three water molecules. The bipyridine complex has two independent moieties in the asymmetric unit related by a non-crystallographic pseudosymmetry center. The thiosulfate anions are quite regular, in contrast with other similar sulfur-coordinated thiosulfates in the literature.

Comment

The thiosulfate group has been shown to be a most versatile ligand, able to coordinate in many different ways, depending both on the characteristics of the cations involved as well as on the environmental conditions. Thus, among the *ca* 50 structures containing the anion which are included in the October 1998 version of the Cambridge Structural Database (CSD; Allen & Kennard, 1993), there are examples where the thiosulfate binds in a mono-, bi- or even tridentate fashion, both in monomeric species as well as in one-, two- or three-dimensional polymers. At the same time, nickel(II) appears to be a rather interesting metal center for thiosulfate complexing because, as a borderline acid ion in the Pearson classification scheme (Pearson, 1973), it would not be expected to exhibit a preference for